

Literatur zum Artikel von Urs Granacher et al.: Posturale Kontrolle im Alter: Effekte von Gleichgewichtstraining, „Praxis Physiotherapie“ 2/2010, S. 131ff.

- (1) Ashton-Miller JA, Wojtys EM, Huston LJ, Fry-Welch D (2001). Can proprioception really be improved by exercises? *Knee Surg Sports Traumatol Arthrosc*, 9, 128-136.
- (2) Banaschewski T, Besmehn F, Zieger H, Rothenberger A (2001). Evaluation of sensorimotor training in children with ADHD. *Percept Mot Skills*, 92, 137-149.
- (3) Beauchet O, Annweiler C, Dubost V, Allali G, Kressig RW, Bridenbaugh S, et al. (2009). Stops walking when talking: a predictor of falls in older adults? *Eur J Neurol*, 16, 786-795.
- (4) Bernier JN, Perrin DH (1998). Effect of coordination training on proprioception of the functionally unstable ankle. *J Orthop Sports Phys Ther*, 27, 264-275.
- (5) Chodzko-Zajko WJ, Proctor DN, Fiatarone Singh MA, Minson CT, Nigg CR, Salem GJ, et al. (2009). American College of Sports Medicine position stand. Exercise and physical activity for older adults. *Med Sci Sports Exerc*, 41, 1510-1530.
- (6) Chong RK, Ambrose A, Carzoli J, Hardison L, Jacobson B (2001). Source of improvement in balance control after a training program for ankle proprioception. *Percept Mot Skills*, 92, 265-272.
- (7) Corden DM, Lippold OC (1996). Age-related impaired reflex sensitivity in a human hand muscle. *J Neurophysiol*, 76, 2701-2706.
- (8) Cumberworth VL, Patel NN, Rogers W, Kenyon GS (2007). The maturation of balance in children. *J Laryngol Otol*, 121, 449-454.
- (9) Earles D, Vardaxis V, Koceja D (2001). Regulation of motor output between young and elderly subjects. *Clin Neurophysiol*, 112, 1273-1279.
- (10) Era P, Sainio P, Koskinen S, Haavisto P, Vaara M, Aromaa A (2006). Postural balance in a random sample of 7,979 subjects aged 30 years and over. *Gerontology*, 52, 204-213.
- (11) Fernie GR, Gryfe CI, Holliday PJ, Llewellyn A (1982). The relationship of postural sway in standing to the incidence of falls in geriatric subjects. *Age & Ageing*, 11, 11-16.
- (12) Gillespie LD, Robertson MC, Gillespie WJ, Lamb SE, Gates S, Cumming RG, et al. (2009). Interventions for preventing falls in older people living in the community. *Cochrane database of systematic reviews (Online)*, CD007146.
- (13) Granacher U (2006). *Neuromuskuläre Leistungsfähigkeit im Alter (> 60 Jahre): Auswirkungen von Kraft- und sensomotorischem Training*. Geislingen/Steige: Maurer Verlag.
- (14) Granacher U, Bergmann S, Gollhofer A (2007). Allgemeine Richtlinien für den Einsatz von sensomotorischem Training im Schulsport. *Sportunterricht*, 9, 259-265.
- (15) Granacher U, Gollhofer A (2007). Die Auswirkungen von sensomotorischem Training im Alter unter besonderer Berücksichtigung alltagsmotorischer Aufgaben. *Med Orthopädieschuhtechnik*, 1, 36-43.
- (16) Granacher U, Gollhofer A (2009). Grundlagen der Bewegungskontrolle und deren funktionelle Bedeutung. *Med Orthopädieschuhtechnik, Sonderheft*, 6-11.
- (17) Granacher U, Gollhofer A, Strass D (2006). Training induced adaptations in characteristics of postural reflexes in elderly men. *Gait & Posture*, 24, 459-466.
- (18) Granacher U, Gruber M, Gollhofer A (2009). Resistance training and neuromuscular performance in seniors. *Int J Sports Med*, 30, 652-657.
- (19) Granacher U, Gruber M, Strass D, Gollhofer A (2007). The impact of sensorimotor training in elderly men on maximal and explosive force production capacity. *Dt Z Sportmed*, 58, 446-451.
- (20) Granacher U, Strass D, Gollhofer A (2008). Effect of aging on power output and

- functional reflex activity. *Med Sci Sports Exerc*, 40, 87-88.
- (21) Granacher U, Zahner L, Gollhofer A (2008). Strength, power, and postural control in seniors: Considerations for functional adaptations and for fall prevention. *Eur J Sport Sci*, 8, 325-340.
- (22) Grey MJ, Ladouceur M, Andersen JB, Nielsen JB, Sinkjaer T (2001). Group II muscle afferents probably contribute to the medium latency soleus stretch reflex during walking in humans. *J Physiol*, 534, 925-933.
- (23) Gruber M, Gollhofer A (2004). Impact of sensorimotor training on the rate of force development and neural activation. *Eur J Appl Physiol*, 92, 98-105.
- (24) Gruber M, Gruber SB, Taube W, Schubert M, Beck SC, Gollhofer A (2007). Differential effects of ballistic versus sensorimotor training on rate of force development and neural activation in humans. *J Strength Cond Res*, 21, 274-282.
- (25) Guimaraes RM, Isaacs B (1980). Characteristics of the gait in old people who fall. *Int Rehabil Med*, 2, 177-180.
- (26) Güllich A, Schmidbleicher D (1999). Struktur der Kraftfähigkeiten und ihrer Trainingsmethoden. *Dt Z Sportmed*, 7/8, 223-234.
- (27) Hausdorff JM, Rios DA, Edelberg HK (2001). Gait variability and fall risk in community-living older adults: a 1-year prospective study. *Arch Phys Med Rehabil*, 82, 1050-1056.
- (28) Heitkamp HC, Horstmann T, Mayer F, Weller J, Dickhuth HH (2001). Gain in strength and muscular balance after balance training. *Int J Sports Med*, 22, 285-290.
- (29) Horak FB, Macpherson JM (1996). Postural orientation and equilibrium. In: Rowell LB, Shepherd JT, editors. *Handbook of Physiology - Section 12: Exercise: Regulation and Integration of Multiple Systems*. New York: Oxford University Press. 255-292.
- (30) Hu MH, Woollacott MH (1994). Multisensory training of standing balance in older adults: II. Kinematic and electromyographic postural responses. *J Gerontol*, 49, M62-M71.
- (31) Hytonen M, Pyykko I, Aalto H, Starck J (1993). Postural control and age. *Acta Otolaryngol*, 113, 119-122.
- (32) Kannus P, Parkkari J, Koskinen S, Niemi S, Palvanen M, Jarvinen M, et al. (1999). Fall-induced injuries and deaths among older adults. *JAMA*, 281, 1895-1899.
- (33) Laessoe U, Hoeck HC, Simonsen O, Voigt M (2008). Residual attentional capacity amongst young and elderly during dual and triple task walking. *Hum Mov Sci*, 27, 496-512.
- (34) Lajoie Y, Teasdale N, Bard C, Fleury M (1993). Attentional demands for static and dynamic equilibrium. *Exp Brain Res*, 97, 139-144.
- (35) Lephart SM, Fu FH (2000). *Proprioception and neuromuscular control in joint stability*. Champaign: Human Kinetics.
- (36) Li F, Harmer P, Fisher KJ, McAuley E, Chaumeton N, Eckstrom E, et al. (2005). Tai Chi and fall reductions in older adults: a randomized controlled trial. *J Gerontol*, 60, 187-194.
- (37) Lin SI, Woollacott MH (2002). Postural muscle responses following changing balance threats in young, stable older, and unstable older adults. *J Mot Behav*, 34, 37-44.
- (38) Liu JX, Eriksson PO, Thornell LE, Pedrosa-Domellof F (2005). Fiber content and myosin heavy chain composition of muscle spindles in aged human biceps brachii. *J Histochem Cytochem*, 53, 445-454.
- (39) Lord SR, Dayhew J (2001). Visual risk factors for falls in older people. *J Am Geriatr Soc*, 49, 508-515.
- (40) Lord SR, Sherrington C, Menz HB (2001). *Falls in older people: Risk factors and strategies for prevention*. Cambridge: Cambridge University Press.
- (41) Lundin-Olsson L, Nyberg L, Gustafson Y (1997). "Stops walking when talking" as a predictor of falls in elderly people. *Lancet*, 349, 617.
- (42) Lutz W (2008). Europe's changing population in a global context. *Eur View*, 7, 237-245.

- (43) Macefield VG, Gandevia SC, Bigland-Ritchie B, Gorman RB, Burke D (1993). The firing rates of human motoneurones voluntarily activated in the absence of muscle afferent feedback. *J Physiol*, 471, 429-443.
- (44) Madureira MM, Takayama L, Gallinaro AL, Caparbo VF, Costa RA, Pereira RM (2007). Balance training program is highly effective in improving functional status and reducing the risk of falls in elderly women with osteoporosis: a randomized controlled trial. *Osteoporos Int*, 18, 419-425.
- (45) Maisonneuve T, Hauw JJ, Dani SU, Hori A, Walter GF (1997). Changes in the peripheral nervous system. In: Dani SU, Hori A, Walter GF, editors. *Principles of neural aging*. Amsterdam: Elsevier. 304-316.
- (46) Maki BE, Cheng KC, Mansfield A, Scovil CY, Perry SD, Peters AL, et al. (2008). Preventing falls in older adults: new interventions to promote more effective change-in-support balance reactions. *J Electromyogr Kinesiol*, 18, 243-254.
- (47) Maki BE, McIlroy WE (1997). The role of limb movements in maintaining upright stance: the "change-in-support" strategy. *Phys Ther*, 77, 488-507.
- (48) Maki BE, McIlroy WE (2005). Change-in-support balance reactions in older persons: an emerging research area of clinical importance. *Neurol Clin*, 23, 751-vii.
- (49) Marchetti GF, Whitney SL (2005). Older adults and balance dysfunction. *Neurol Clin*, 23, 785-805, vii.
- (50) McNeil CJ, Doherty TJ, Stashuk DW, Rice CL (2005). Motor unit number estimates in the tibialis anterior muscle of young, old, and very old men. *Muscle & Nerve*, 31, 461-467.
- (51) Meunier S, Pierrot-Deseilligny E (1989). Gating of the afferent volley of the monosynaptic stretch reflex during movement in man. *J Physiol*, 419, 753-763.
- (52) Minetti AE (2000). The three modes of terrestrial locomotion. In: Nigg BM, MacIntosh BR, Mester J, editors. *Biomechanics and Biology of Movement*. Champaign: Human Kinetics. 67-78.
- (53) Mion LC, Gregor S, Buettner M, Chwirchak D, Lee O, Paras W (1989). Falls in the rehabilitation setting: incidence and characteristics. *Rehabil Nurs*, 14, 17-22.
- (54) Miwa T, Miwa Y, Kanda K (1995). Dynamic and static sensitivities of muscle spindle primary endings in aged rats to ramp stretch. *Neurosci Lett*, 201, 179-182.
- (55) Morisawa Y (1998). Morphological study of mechanoreceptors on the coracoacromial ligament. *J Orthop Sci*, 3, 102-110.
- (56) Mynark RG, Koceja DM (2001). Effects of age on the spinal stretch reflex. *J Appl Biomech*, 17, 188-203.
- (57) Mynark RG, Koceja DM (2002). Down training of the elderly soleus H reflex with the use of a spinally induced balance perturbation. *J Appl Physiol*, 93, 127-133.
- (58) Paterno MV, Myer GD, Ford KR, Hewett TE (2004). Neuromuscular training improves single-limb stability in young female athletes. *J Orthop Sports Phys Ther*, 34, 305-316.
- (59) Pavol MJ, Owings TM, Foley KT, Grabiner MD (2001). Mechanisms leading to a fall from an induced trip in healthy older adults. *J Gerontol*, 56, M428-M437.
- (60) Pellecchia GL (2005). Dual-task training reduces impact of cognitive task on postural sway. *J Mot Behav*, 37, 239-246.
- (61) Rubenstein LZ (2006). Falls in older people: epidemiology, risk factors and strategies for prevention. *Age & Ageing*, 35 Suppl 2, ii37-ii41.
- (62) Scaglioni G, Narici MV, Maffiuletti NA, Pensini M, Martin A (2003). Effect of ageing on the electrical and mechanical properties of human soleus motor units activated by the H reflex and M wave. *J Physiol*, 548, 649-661.
- (63) Shaffer SW, Harrison AL (2007). Aging of the somatosensory system: a translational perspective. *Phys Ther*, 87, 193-207.
- (64) Shumway-Cook A, Woollacott M (2001). *Motor Control: Theory and Practical Applications*. Philadelphia: Lippincott Williams & Wilkins.
- (65) Silsupadol P, Lugade V, Shumway-Cook A, van Donkelaar P, Chou LS, Mayr U, et al. (2009). Training-related changes in dual-task walking performance of elderly persons with balance impairment: A double-blind, randomized controlled trial. *Gait*

& Posture, 29, 634-639.

- (66) Silsupadol P, Shumway-Cook A, Lugade V, van Donkelaar P, Chou LS, Mayr U, et al. (2009). Effects of single-task versus dual-task training on balance performance in older adults: a double-blind, randomized controlled trial. *Arch Phys Med Rehabil*, 90, 381-387.
- (67) Steadman J, Donaldson N, Kalra L (2003). A randomized controlled trial of an enhanced balance training program to improve mobility and reduce falls in elderly patients. *J Am Geriatr Soc*, 51, 847-852.
- (68) Stevens JA, Corso PS, Finkelstein EA, Miller TR (2006). The costs of fatal and non-fatal falls among older adults. *Inj Prev*, 12, 290-295.
- (69) Swash M, Fox KP (1972). The effect of age on human skeletal muscle. Studies of the morphology and innervation of muscle spindles. *J Neurol Sci*, 16, 417-432.
- (70) Tang PF, Woollacott MH (1998). Inefficient postural responses to unexpected slips during walking in older adults. *J Gerontol A Biol Sci Med Sci*, 53, M471-M480.
- (71) Taube W, Gruber M, Beck S, Faist M, Gollhofer A, Schubert M (2007). Cortical and spinal adaptations induced by balance training: correlation between stance stability and corticospinal activation. *Acta Physiol (Oxf)*, 189, 347-358.
- (72) Taube W, Gruber M, Gollhofer A (2008). Spinal and supraspinal adaptations associated with balance training and their functional relevance. *Acta Physiol (Oxf)*, 193, 101-116.
- (73) Terao S, Sobue G, Hashizume Y, Li M, Inagaki T, Mitsuma T (1996). Age-related changes in human spinal ventral horn cells with special reference to the loss of small neurons in the intermediate zone: a quantitative analysis. *Acta Neuropathol (Berl)*, 92, 109-114.
- (74) Vandervoort AA (2002). Aging of the human neuromuscular system. *Muscle & Nerve*, 25, 17-25.
- (75) Williams HG, Burke JR, McClenaghan BA, Hirth V, Huber G (1997). Balance control: mechanisms of adaptation to sensory-motor integration training in the elderly. Proceedings of the IV International Conference on Physical Activity and Aging. Heidelberg: Health Promotion Publications. 118-130.
- (76) Wolf SL, Barnhart HX, Ellison GL, Coogler CE (1997). The effect of Tai Chi Quan and computerized balance training on postural stability in older subjects. Atlanta FICSIT Group. Frailty and Injuries: Cooperative Studies on Intervention Techniques. *Phys Ther*, 77, 371-381.
- (77) Wolf SL, Barnhart HX, Kutner NG, McNeely E, Coogler C, Xu T (1996). Reducing frailty and falls in older persons: an investigation of Tai Chi and computerized balance training. Atlanta FICSIT Group. Frailty and Injuries: Cooperative Studies of Intervention Techniques. *J Am Geriatr Soc*, 44, 489-497.
- (78) Woollacott M, Shumway-Cook A (2002). Attention and the control of posture and gait: a review of an emerging area of research. *Gait & Posture*, 16, 1-14.
- (79) Wuker N, Rudert M (1999). Lateral ankle ligament rupture. When is surgical management indicated and when conservative therapy preferred? *Orthopade*, 28, 476-482.