

**Literatur zum Artikel von Holm Thieme: Sturzprävention in der Physiotherapie,
„Praxis Physiotherapie“ 1/2010, S. 60ff.**

Balash Y, Peretz C, Leibovich G, Herman T, Hausdorff JM, Giladi N. Falls in outpatients with Parkinson's disease: frequency, impact and identifying factors. *J Neurol*. 2005;252(11):1310-5.

Barak Y, Wagenaar RC, Holt KG. Gait characteristics of elderly people with a history of falls: a dynamic approach. *Phys Ther*. 2006;86(11):1501-10.

Berg K, Wood-Dauphinee S, Gayton D. Measuring balance in the elderly: preliminary development of an instrument. *Physiother Can*. 1989;41:304-10.

Bloem BR, Boers I, Cramer M, Westendorp RG, Gerschlager W. Falls in the elderly. I. Identification of risk factors. *Wien Klin Wochenschr* 2001;113(10):352-62.

Bogle Thorbahn LD, Newton RA. Use of the Berg Balance Test to predict falls in elderly persons. *Phys Ther* 1996;76(6):576-83.

Chan BK, Marshall LM, Winters KM, Faulkner KA, Schwartz AV, Orwoll ES. Incident fall risk and physical activity and physical performance among older men: the Osteoporotic Fractures in Men Study. *Am J Epidemiol*. 2007;165(6):696-703.

Chang JT, Morton SC, Rubenstein LZ, Mojica WA, Maglione M, Suttorp MJ, Roth EA, Shekelle PG. Interventions for the prevention of falls in older adults: systematic review and meta-analysis of randomised clinical trials. *BMJ*. 2004;328(7441):680.

Clemson L, Cumming RG, Roland M. Case-control study of hazards in the home and risk of falls and hip fractures. *Age Ageing*. 1996;25(2):97-101.

Carroll NV, Slattum PW, Cox FM. The cost of falls among the community-dwelling elderly. *J Manag Care Pharm*. 2005;11(4):307-16.

Carter SE, Campbell EM, Sanson-Fisher RW, Redman S, Gillespie WJ. Environmental hazards in the homes of older people. *Age Ageing* 1997;26(3):195-202.

Faber MJ, Bosscher RJ, Chin A Paw MJ, van Wieringen PC. Effects of exercise programs on falls and mobility in frail and pre-frail older adults: A multicenter randomized controlled trial. *Arch Phys Med Rehabil*. 2006;87(7):885-96.

Gates S, Smith LA, Fisher JD, Lamb SE. Systematic review of accuracy of screening instruments for predicting fall risk among independently living older adults. *J Rehabil Res Dev.* 2008;45(8):1105-16.

Giampaoli S, Ferrucci L, Cecchi F, Lo Noce C, Poce A, Dima F, Santaquilani A, Vescio MF, Menotti A. Hand-grip strength predicts incident disability in non-disabled older men. *Age Ageing.* 1999;28(3):283-8.

Gillespie LD, Robertson MC, Gillespie WJ, Lamb SE, Gates S, Cumming RG, Rowe BH. Interventions for preventing falls in older people living in the community. *Cochrane Database Syst Rev.* 2009;(2):CD007146.

Gillespie LD, Gillespie WJ, Robertson MC, Lamb SE, Cumming RG, Rowe BH. Interventions for preventing falls in elderly people. *Cochrane Database Syst Rev.* 2003;(4):CD000340.

Heinze C, Lahmann N, Dassen T. Sturzfrequenz in deutschen Krankenhäusern. *Gesundheitswesen.* 2002;64(11):598-601.

Lundin-Olsson L, Nyberg L, Gustafson Y. "Stops walking when talking" as a predictor of falls in elderly people. *Lancet.* 1997;349(9052):617.

Luukinen H, Koski K, Honkanen R, Kivela SL. Incidence of injury-causing falls among older adults by place of residence: a population-based study. *J Am Geriatr Soc* 1995;43(8):871-6.

Mackintosh SF, Goldie P, Hill K. Falls incidence and factors associated with falling in older, community-dwelling, chronic stroke survivors (> 1 year after stroke) and matched controls. *Aging Clin Exp Res.* 2005;17(2):74-81.

Marks D. Assessment: Tinetti Test. *Physiopraxis* 2006;2.

Parker MJ, Gillespie LD, Gillespie WJ. Hip protectors for preventing hip fractures in the elderly. *Cochrane Database Syst Rev.* 2004;(3):CD001255.

Parkkari J, Heikkila J, Kannus IP. Acceptability and compliance with wearing energy-shunting hip protectors: a 6-month prospective follow-up in a Finnish nursing home. *Age Ageing.* 1998;27(2):225-9.

Petridou ET, Manti EG, Ntinapogias AG, Negri E, Szczerbinska K. What works better for community-dwelling older people at risk to fall?: a meta-analysis of multifactorial versus physical exercise-alone interventions. J Aging Health. 2009;21(5):713-29.

Pientka L, Friedrich C. Die Kosten hüftgelenksnaher Frakturen in Deutschland: Eine prospektive Untersuchung. Zeitschrift Gerontol Geriatr. 1999;32: 326-332

Pluijm SM, Smit JH, Tromp EA, Stel VS, Deeg DJ, Bouter LM, Lips P. A risk profile for identifying community-dwelling elderly with a high risk of recurrent falling: results of a 3-year prospective study. Osteoporos Int 2006;17(3):417-25.

Scherfer E, Bohls C, Freiberger E, Heise KF, Hogan D. Berg Balance Scale – deutsche Version. Übersetzung eines Instruments zur Beurteilung von Gleichgewicht und Sturzgefährdung. Physioscience 2006;2:59-66.

Shumway-Cook A, Cioc MA, Gruber W, Robinson C. Incidence of and risk factors for falls following hip fracture in community-dwelling older adults. Phys Ther. 2005;85(7):648-55.

Shumway-Cook A, Brauer S, Woollacott M. Predicting the probability for falls in community-dwelling older adults using the Timed Up & Go Test. Phys Ther. 2000;80(9):896-903.

Shumway-Cook A, Baldwin M, Polissar NL, Gruber W. Predicting the probability for falls in community-dwelling older adults. Phys Ther 1997;77(8):812-9.

Spreng T, Spreng K, Gulich M, et al. Stürze älterer Menschen: Ergebnisse einer epidemiologischen Erhebung. ZfA 2000;76 (9).

Tinetti ME. Performance-Oriented assessment of mobility problems. J Am Geriatr Soc 1986;34:119-26.

Vellas BJ, Wayne SJ, Romero LJ, Baumgartner RN, Garry PJ. Fear of falling and restriction of mobility in elderly fallers. Age Ageing 1997;26(3):189-93.

Villar MT, Hill P, Inskip H, Thompson P, Cooper C. Will elderly rest home residents wear hip protectors? Age Ageing. 1998;27(2):195-8.

Voukelatos A, Cumming RG, Lord SR, Rissel C. A randomized, controlled trial of tai chi for the prevention of falls: the Central Sydney tai chi trial.

J Am Geriatr Soc. 2007;55(8):1185-91.

Zeitler H, Gulich M. Ältere Sturzpatienten. Leitlinie der AWMF. (<http://www.uni-duesseldorf.de/AWMF/II/053-004.htm>, Zugang: 08.11.2007)